



SYSTEM OF NARCOTICS IDENTIFICATION

INSTRUCTIONS FOR USE

TEST K

FOR: Heroin, Codeine, Morphine

Test Function

1. This reagent presumptively identifies Heroin, Codeine and Morphine.

How to Use Test K

1. Remove clip and open test container.
2. Insert an amount of powdered suspect material that would fit inside this circle into the test pouch. Reseal with clip and tap gently to assure material falls to bottom of pouch.
3. With printed side of test pouch facing you, break ampoule by squeezing the center of the ampoule with the tip of the thumb and forefinger. Agitate gently and observe color or color changes.

Interpretation of Resulting Colors

1. An immediate green color, changing to purple within 2 or 3 minutes indicates the presence of Heroin.
2. An immediate blue-green color, changing to a gray color within 2 or 3 minutes indicates the presence of Morphine.
3. An immediate stable blue color indicates the presence of Codeine.
4. It is advisable that this test be used in conjunction with others in the Polytesting System.

CAUTION

- A. Since these tests contain strong acids, it is suggested that Test F - Acid Neutralizer be used after testing and before disposal.
- B. Before discarding used test pouches, remove clip and add one measure of acid neutralizer from Test F.
- C. Add slowly to prevent bubbling over.
- D. DO NOT reseal the test pouch until effervescence has completely stopped.
- E. Reseal pouch with clip and discard in a tamper-free disposal unit.
- F. No attempt should be made to crush the glass particles after the ampoules are broken.
- G. **Antidote:** In case of contact, immediately flush eyes or skin with water for at least 15 minutes while removing contaminated clothing and shoes. If swallowed, do not give emetics. Contact a physician.

WARNING

These tests are NOT designed for use with liquid samples.



13386 International Parkway
Jacksonville, Florida 32218
904.485.1836 FAX: 904.741.5407
© 2105 Safariland, LLC

A Brand of the Safariland Group